

Other Pertinent Reading

Newer Ionization Methods

42. Karas M, Bachman D, Bahr U, & Hillenkamp F. Matrix-assisted ultraviolet laser desorption of non-volatile compounds. *Int. J. Mass Spectrom. Ion Process* 78: 53 (1987).
43. Beavis RC & Chait BT. Matrix assisted laser desorption ionization mass spectrometry of proteins. *Methods Enzymol.* 270: 519 (1996).
44. Fenn JB. Electrospray Wings for Molecular Elephants (Nobel Lecture). *Angew. Chem. Int. Ed.* 42, 3871 – 3894 (2003).

Mass Analyzers

45. Paul W. Electromagnetic traps for charged and neutral particles. *Reviews of Modern Physics* 62: 531-540 (1990).
46. March RE. Quadrupole ion trap mass spectrometry: A view at the turn of the century. *Int. J. Mass Spectrom.* 200: 285-312 (2000).
47. Marshall AG, Hendrickson CL, & Jackson GS. Fourier Transform Ion Cyclotron Resonance Mass Spectrometry: A Primer. *Mass Spectrometry Reviews* 17: 1-35 (1998).
48. Marshall AG. Milestones in Fourier transform ion cyclotron resonance mass spectrometry development *Int. J. Mass Spectrom. Ion Processes* 200: 331-356 (2000).
49. Hu Q, Noll RJ, Li H, Makarov A, Hardman M & Cooks RG. The Orbitrap: a new mass spectrometer. *J. Mass Spectrom.* 40: 430-443 (2005).

Newer Tandem Mass Spectrometers

50. Chernushevich IV, Loboda AV, & Thomson BA. An introduction to quadrupole/time-of-flight mass spectrometry *J. Mass Spectrom.* 36: 849-865 (2001). [Describes the QTOF]
51. Schwartz JC, MW, & Syka JEP. A Two-Dimensional Quadrupole Ion Trap Mass Spectrometer. *J. Am. Soc. Mass. Spectrom.* 13, 659 – 669 (2002). [Describes the LTQ]
52. Londry FA & Hager JW. Mass selective axial ion ejection from a linear quadrupole ion trap. *J. Am. Soc. Mass Spectrom.* 14: 1130-1147 (2003). [Describes the QTRAP]

Isotope Dilution GC/MS (EI) to Quantitate and Confirm Drugs of Abuse

53. Bodor G, Roggeman R, & Turk J: Variations in abundance of the molecular ion of the derivatized cocaine metabolite benzoylecgonine. *Clin. Chem.* 36: 742-747 (1990).

Negative Ion Electron Capture Measurement of Bioactive Substances

54. Leis HJ, Fauler G, Rechberger GN, & Windischhofer W. Electron-Capture Mass Spectrometry: A Powerful Tool in Biomedical Trace Level Analysis. *Current Medicinal Chemistry* 11: 1585-1594 (2004).
55. Stickle D, Bohrer A, Berger R, Morrisey J, Klahr S, & Turk J. Quantitation of the putative neurotransmitter agmatine as the hexafluoroacetylacetone derivative by stable isotope dilution gas chromatography and negative ion chemical ionization mass spectrometry. *Anal. Biochem.* 238: 129-136 (1996).
56. Kempe K, Hsu FF, Bohrer A, & Turk J. Isotope dilution mass spectrometric measurements indicate that arachidonylethanolamide, the proposed endogenous ligand of the cannabinoid receptor, accumulates in rat brain tissue post-mortem but is contained at low levels in or is absent from fresh tissue. *J. Biol. Chem.* 271: 17287-17295 (1996).
57. Peters FT, Samyn N, Kraemer T, Riedel WJ, Maurer HH. Negative-Ion Chemical Ionization Gas Chromatography–Mass Spectrometry Assay for Enantioselective Measurement of Amphetamines in Oral Fluid: Application to a Controlled Study with MDMA and Driving Under the Influence Cases. *Clinical Chemistry* 53: 702–710 (2007).

Quantitative Linked Tandem MS Scanning Modes on Triple Stage Quadrupoles: Therapeutic Drug Level Monitoring and Biochemical Measurements

58. Taylor PJ, Jones A, Balderson GA, Lynch SV, Norris RLG, & Pond SM Sensitive, specific quantitative analysis of tacrolimus (FK506) in blood by liquid chromatography-electrospray tandem mass spectrometry. *Clinical Chemistry* 42: 279-285 (1996).
59. Ramanadham S, Hsu FF, Zhang S, Jin C, Bohrer A, Ma Z, & Turk J. Involvement of the Group VIA Phospholipase A₂ (iPLA₂β) in endoplasmic reticulum stress-induced apoptosis in insulinoma cells. *Biochemistry* 43: 918-930 (2004).

Peptide/Protein Characterization by MALDI/TOF/MS and LC/ESI/MS/MS

60. Le Blanc JCY, Hager JW, Ilisiu AMP, Hunter C, Zhong F, Chu I. Unique scanning capabilities of a new hybrid linear ion trap mass spectrometer (Q TRAP) used for high sensitivity proteomics applications. *Proteomics* 3: 859-869 859 (2003).
61. Ramanadham S, Song H, Hsu FF, Zhang S, Crankshaw M, Grant G, Newgard C, & Turk J. Pancreatic islets and insulinoma cells express a novel isoform of Group VIA PLA₂ (iPLA₂β) that participates in glucose-stimulated insulin secretion and is not produced by alternate splicing of the iPLA₂β transcript. *Biochemistry* 42: 13929-40 (2003).
62. Hortin GL. The MALDI-TOF Mass Spectrometric View of the Plasma Proteome and Peptidome. *Clinical Chemistry* 52: 1223-1237 (2006).
63. Albrethsen J. Reproducibility in Protein Profiling by MALDI-TOF Mass Spectrometry *Clinical Chemistry* 53: 852- 858 (2007).

MALDI/TOF/MS Methods for Genotyping

64. Jalanko A, Kere J, Savilahti E, Schwartz M, Syvflnen AC, Ranki M, & Soderlund H. Screening for Defined Cystic Fibrosis Mutations by Solid-Phase Minisequencing. *Clin. Chem.* 38: 39-43 (1992).
65. Nikiforov TT, Rendle RB, Goelet P, Rogers YH, Kotewicz ML, Anderson S, Trainor GL, & Knapp MR. Genetic Bit Analysis: a solid phase method for typing single nucleotide polymorphisms. *Nucleic Acids Research* 22: 4167-4175 (1994).
66. Braun A, Little DP, & Koster H. Detecting CFTR gene mutations by using primer oligo base extension and mass spectrometry. *Clinical Chemistry* 43: 1151-1158 (1997).
67. Ross P, Hall L, Smirnov I, & Haff L. High level multiplex genotyping by MALDI-TOF mass spectrometry. *Nature Biotechnology* 16: 1347-1351 (1998).
68. Kepper P, Reinhardt R, Dahl A, Lehrach H, & Sauer S. Matrix-Assisted Laser Desorption/Ionization Mass Spectrometric Analysis of DNA on Microarrays. *Clinical Chemistry* 52: 1303-1310 (2006).
69. Kao HY, Su YN, Liao HK, MS, & Yu-Ju Chen YJ. Determination of SMN1/SMN2 Gene Dosage by a Quantitative Genotyping Platform Combining Capillary Electrophoresis and MALDI-TOF Mass Spectrometry. *Clin. Chem.* 52: 361-369 (2006).
70. Blievernicht JK, Schaeffeler E, Klein K, Eichelbaum M, Schwab M, & Zanger UM. MALDI-TOF Mass Spectrometry for Multiplex Genotyping of CYP2B6 Single-Nucleotide Polymorphisms. *Clinical Chemistry* 53: 24-33 (2007).

Drug Screening with 3D Ion Trap Tandem MS Systems

71. Fitzgerald RL, Rivera JD, & Herold DA. Broad Spectrum Drug Identification Directly from Urine, Using Liquid Chromatography-Tandem Mass Spectrometry. *Clinical Chemistry* 45: 1224 -1234 (1999). [Drug screening with an LCQ].

Drug Screening with Triple Stage Quadrupole (TSQ) Tandem MS Systems

72. Weinmann W, Wiedemann A, Eppinger B, Renz M, & Svoboda M. Screening for Drugs in Serum by Electrospray Ionization/Collision-Induced Dissociation and Library Searching. *J. Am. Soc. Mass Spectrom.* 10: 1028 –1037 (1999).
73. Weinmann W, Stoertzel M, Vogt S, Svoboda M, & Schreiber A. Tuning compounds for electrospray ionization/in-source collision-induced dissociation and mass spectra library searching. *J. Mass Spectrom.* 36: 1013 – 1023 (2001).
74. Muller C, Schafer P, Stortzel M, Vogt S, & Weinmann W. Ion suppression effects in liquid chromatography–electrospray-ionization transport-region collision induced dissociation mass spectrometry with different serum extraction methods for systematic toxicological analysis with mass spectra libraries. *J. Chromatogr. B* 773: 47-52 (2002).
75. Gergov M, Ojanpera I, & Vuori E. Simultaneous screening for 238 drugs in blood by liquid chromatography–ionspray tandem mass spectrometry with multiple-reaction monitoring. *J. Chromatogr. B* 795: 41-53 (2003).
76. Maralikova B & Weinmann W. Confirmatory analysis for drugs of abuse in plasma and urine by high-performance liquid chromatography–tandem mass spectrometry with respect to criteria for compound identification. *J. Chromatogr. B* 811: 21-30 (2004).
77. Quintela O, Sauvage FL, Charvier F, Gaulier JM, Lachatre G, Marquet P. Liquid Chromatography–Tandem Mass Spectrometry for Detection of Low Concentrations of 21 Benzodiazepines, Metabolites, and Analogs in Urine: Method with Forensic Applications. *Clin. Chem.* 52: 1346 –1355 (2006).
78. Øiestad EL, Johansen U, & Christophersen AS. Drug Screening of Preserved Oral Fluid by Liquid Chromatography-Tandem Mass Spectrometry. *Clin. Chem.* 53:300-309 (2007).

Drug Screening with QTRAP Hybrid Quadrupole/Linear Ion Trap Tandem MS Systems

79. Marquet P, Saint-Marcoux F, Gamble TN, Leblanc JCY. Comparison of a preliminary procedure for the general unknown screening of drugs and toxic compounds using a quadrupole-linear ion-trap mass spectrometer with a liquid chromatography–mass spectrometry reference technique. *J. Chromatogr. B* 789: 9–18 (2003).
80. Herrin GL, McCurdy HH, & Wall WH. Investigation of an LC–MS–MS (QTrap) Method for the Rapid Screening and Identification of Drugs in Post mortem Toxicology Whole Blood Samples. *Journal of Analytical Toxicology* 29: 599-607 (2005).
81. Mueller CA, Weinmann W, Dresen S, Schreiber A, & Gergov M. Development of a multi-target screening analysis for 301 drugs using a QTrap liquid chromatography/tandem mass spectrometry system and automated library searching. *Rapid Commun. Mass Spectrom.* 19: 1332-1338 (2005).
82. Jansen R, Lachatre G, & Marquet P. LC-MS/MS systematic toxicological analysis: Comparison of MS/MS spectra obtained with different instruments and settings. *Clinical Biochemistry* 38: 362 – 372 (2005). [Compares drug screening on QTRAP vs. TSQ]
83. Sauvage FL, Saint-marcoux F, Duretz B, Deporte D, Lachatre G, & Marquet P Screening of Drugs and Toxic Compounds with Liquid Chromatography-Linear Ion Trap Tandem Mass Spectrometry. *Clin. Chem.* 52: 1735–1742 (2006).

Drug Screening with QTOF Hybrid Quadrupole/Time-of-Flight Tandem MS Systems

84. Decaestecker TN, Clauwaert KM, Van Bocxlaer JF, Lambert WE, Van den Eeckhout EG, Van Peteghem CH, De Leenheer AP. Evaluation of automated single mass spectrometry to tandem mass spectrometry function switching for comprehensive drug profiling analysis using a quadrupole time-of-flight mass spectrometer. *Rapid Commun. Mass Spectrom.* 14: 1787-92 (2000).
85. Decaestecker TN, Van de Castele SR, Wallemacq PE, Van Peteghem, CH Defore DL & Van Bocxlaer JF. Information-Dependent Acquisition-Mediated LC-MS/MS Screening Procedure with Semiquantitative Potential. *Anal. Chem.* 76: 6365-6373 (2004).

Accurate Mass Measurements by TOF/MS and FTMS in Drug Screening

86. Ojanperä I, Pelander A, Laks S, Gergov M, Vuori E, & Witt M. Application of Accurate Mass Measurement to Urine Drug Screening. *Journal of Analytical Toxicology* 29: 34-40 (2005).